

small black fruits of *Murraya koenigii* (Curry Leaf Tree; Fig. 1B), that measured ca. 4–5 mm in diameter. This lizard was 3 m up in the tree and we watched it feed for 10 min as it ate at least on four fruits.

To our knowledge, this is the first documented record of fruits in the diet of *F. pardalis*. Lizard frugivory is often considered an “insular phenomenon” and has been rarely reported in Chamaeleonidae (Valido and Olesen 2019. *Front. Ecol. Evol.* 7:49). The only other two chameleon species reported to eat fruit that we are aware of are *Chamaeleo chamaeleon* (Meiri 2018. *Global Ecol. Biogeogr.* 27:1168–1172) and *Furcifer oustaleti* (Takahashi 2008. *J. Herpetol.* 42:760–763). It remains unclear how common frugivory is in chameleons and further work is needed.

We are grateful to the NGO Nature Océan Indien for support.

**MICKAËL SANCHEZ**, Nature Océan Indien, 97429 Petite Île, Reunion Island, France (e-mail: mickael.sanchez974@gmail.com); **LAURENT DEBORDES**, 11 rue du Marais, 79510 Coulon, France (e-mail: 79delau@gmail.com); **ANTOINE BAGLAN**, 320 TER D, Route de Bois de Nêfles, 97490 Sainte Clotilde, Reunion Island, France (e-mail: antoine\_baglan@hotmail.com).

#### **HELODERMA SUSPECTUM (Gila Monster). HIBERNATION.**

During hibernation, *Heloderma suspectum* remain in their overwintering shelters (commonly crevices in rocky slopes or boulders, or burrows of tortoises or rodents) for several months (Beck 2005. *Biology of Gila Monsters and Beaded Lizards*. University of California Press, Berkeley, California. 211 pp.). The duration of hibernation for *H. suspectum* varies by population, individual, and age class. Beck (2005, *op. cit.*) suggests that individuals remain within shelters for between 75 and 120 d (Beck 2005, *op. cit.*). Gienger (2003. *Natural History of the Gila Monster in Nevada*. M.S. Thesis, University of Nevada, Reno, Reno, Nevada. 55 pp.) reported an average hibernation period for *H. suspectum* in southern Nevada of 130 d, with the longest reported hibernation lasting 154 d. Here, we report on an individual whose hibernation period far exceeded any previously reported case.

During a long-term radio telemetry study at a Mojave Desert site in southern Nevada, USA (near Las Vegas), we observed an individual *H. suspectum* overwinter for 232 d, 78 d longer than the maximum duration previously documented. This individual was tracked to its overwintering shelter (1357 m elev.) on 11 November 2020, where it remained until 1 July 2021. Location was confirmed once per week in the field using radio telemetry from August–March, and then daily during the active season (April–July). On its most recent capture on 10 May 2020 this lizard had an SVL of 266 mm, mass of 315 g, and tail volume of 40 mL and was categorized as a sub-adult of unknown sex based on its size relative to breeding adults in the same population.

This nearly 8-mo residency within the hibernaculum suggests the potential for individual *H. suspectum* to remain inactive for a substantially longer period into the activity season than previously thought. This extended quiescent period corresponded with a 238-d period of exceptional drought at the field site, lasting from 1 December 2020 until 27 July 2021 (www.drought.gov; 1 Sept 2021). For individuals active above-ground, particularly during drought years when prey availability is greatly reduced, the energetic return on foraging activities may not outweigh the energetic costs of leaving an overwintering shelter. This may suggest that some *H. suspectum* following an optimal foraging strategy extend their hibernation period into the active season as a means of energy conservation.

**CONNOR J. HUGHES** (e-mail: connorhughes926@gmail.com) and

**JOCELYN B. STALKER**, Center of Excellence for Field Biology, Department of Biology, Austin Peay State University, Clarksville, Tennessee 37040, USA (e-mail: jocelynstalker@gmail.com); **JASON L. JONES**, Nevada Department of Wildlife, Las Vegas, Nevada 89120, USA (e-mail: jljones@ndow.org); **C. M. GIENGER**, Center of Excellence for Field Biology, Department of Biology, Austin Peay State University, Clarksville, Tennessee 37040, USA (e-mail: giengerc@apsu.edu).

#### **HOLCOSUS FESTIVUS (Central American Whiptail Lizard).**

**NUPTIAL GIFT.** Signaling the ability to provision limited resources to a mate’s reproductive success should be favored by sexual selection (Andersson 1994. *Sexual Selection*. Princeton University Press, Princeton, New Jersey. 624 pp.; Wedell and Ritchie 2004. *Anim. Behav.* 67:1059–1065). In many taxa, female mate choice is driven by courtship offerings in the form of nutritional items, commonly known as nuptial gifts where males present their partners with feeding gifts before, during, or after courtship and/or copulating (Lewis and South 2012. *Adv. Study Behav.* 44:53–97). Such mating effort has been widely reported across invertebrate species, but it has been less reported among vertebrates, especially reptiles (Lewis and South 2012, *op. cit.*). Here, we report on a novel mating ritual in the lizard *Holcosus festivus* from a tropical lowland rainforest in Costa Rica.

On 22 August 2020, at ca. 1233 h, we observed an adult male *H. festivus* offer a nuptial gift to a female at Veragua Rainforest Preserve, in Las Brisas de Veragua, Limón, Costa Rica (9.9264°N, 83.1876°W; WGS 84; 420 m elev.). The nuptial gift was an adult *Craugastor fitzingeri* (Fitzinger’s Rain Frog), which the male lizard held in its mouth and had already killed. After ca. 1 min, the male approached the receptive female, which was stationary and remained so, before the male positioned its prey in front of her at a distance of ca. 40 cm. Subsequently, the male circled the female once every minute for 4 min while also performing vertical head movements and tail vibrations in front of her at a distance of ca. 50 cm. Afterwards, the male dropped the frog ca. 10 cm from the female, and she picked it up and consumed it vent first. Fifteen seconds after the female started eating the frog, the male positioned himself on top of her in diagonal position, held onto the female’s back, and wrapped his tail under hers, vent-to-vent, and they mated for about 2 min (Fig. 1). Approximately 3 min after




FIG. 1. *Holcosus festivus* breeding in Veragua Rainforest, Costa Rica. Note the female (bottom) eating a frog that was offered by the male (top) as a nuptial gift before copulation.